

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 1972		2. REPORT TYPE		3. DATES COVERED 00-00-1972 to 00-00-1972	
4. TITLE AND SUBTITLE Description of the Egg of Aedes (Diceromyia) furcifer (Edwards) (Diptera: Culicidae)				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Walter Reed Army Institute of Research, Department of Entomology, Washington, DC, 20012				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 3	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Description of the Egg of Aedes (Diceromyia) furcifer
(Edwards) (Diptera: Culicidae)¹

John F. Reinert²
Department of Entomology
Walter Reed Army Institute of Research
Walter Reed Army Medical Center
Washington, D. C. 20012

The eggs of Aedes furcifer (Edwards) are described and illustrated for the first time herein. This is also the first description of the eggs for any species of the subgenus Diceromyia Theobald. The following description is based on 4 mature eggs extracted from the abdomen of a museum specimen which had the following collection data on the labels: UGANDA, Bwamba Province, April 1948, E. C. C. van Someren collector, and SEAMP Accession Number 111. The nomenclature used follows that of Kalpage and Brust (1968) and Craig and Horsfall (1960).

Eggs of furcifer are very resistant to desiccation as reported by Muspratt (1955: 174). He obtained hatching when he flooded eggs that had been laid in bamboo pots more than 14 months earlier. This drought-resistance of the eggs is an adaptation that enables the species, and probably, other members of the subgenus, to survive dry periods. The only other information on the eggs of Diceromyia is given by Mattingly (1959: 2, 43) for the Malayan species franciscoi Mattingly. He reports eggs of this species were collected from bamboo pots hung about 30 feet from the ground in a mango tree.

DESCRIPTION OF THE EGG
(Figs. 1-2)

Shape (Fig. 1). Broadly fusiform, with ends bluntly rounded; anterior end with sharp taper, posterior end with gradual taper, greatest diameter between anterior third and middle. Size. Length 543-545 microns; width at widest point 182-183 microns. Color. Dark brown. Chorion (Fig. 2). Reticulation uniform, composed of pentagonal and hexagonal cells, and occasionally diamond-shaped ones, cell walls raised and prominent.

¹This work was supported by Research Contract No. DA-49-193-MD-1672 from the U. S. Army Medical Research and Development Command, Office of the Surgeon General, and carried out at the Southeast Asia Mosquito Project, Smithsonian Institution, Washington, D. C. 20560.

²Major, Medical Service Corps, U. S. Army.

DISCUSSION

The reticulation of furcifer closely resembles that of Aedes (Ochlerotatus) campestris Dyar and Knab from the semi-arid plains of North America and Aedes (Ochlerotatus) fitchii (Felt and Young) from the forested areas of the northern Nearctic Region.

Eggs of furcifer have a resemblance in shape and chorion sculpturing to those of species in the subgenus Ochlerotatus Lynch Arribalzaga and show little similarity to those of the known species of Aedimorphus Theobald and Stegomyia Theobald.

ACKNOWLEDGEMENTS

I am grateful to Dr. Botha de Meillon, Principle Investigator, Southeast Asia Mosquito Project (SEAMP), and LTC Bruce F. Eldridge, Chief of the Department of Entomology, Walter Reed Army Institute of Research, for reviewing the manuscript. Thanks are given to Sheila Ford, SEAMP, for preparing the illustrations. I am especially appreciative to my wife, Mollie, for typing the manuscript.

LITERATURE CITED

- Craig, G. B., Jr., and W. R. Horsfall. 1960. Eggs of flood-water mosquitoes. VII. Species of Aedes common in the southeastern United States (Diptera: Culicidae). Ann. ent. Soc. Am. 53: 11-18.
- Kalpage, K. S., and R. A. Brust. 1968. Mosquitoes of Manitoba. I. Descriptions and a key to Aedes eggs (Diptera: Culicidae). Can. J. Zool. 46: 699-718.
- Mattingly, P. F. 1959. The culicine mosquitoes of the Indo-malayan area. Part IV. Genus Aedes Meigen, subgenera Skusea Theobald, Diceromyia Theobald, Geoskusea Edwards and Christophersiomyia Barraud. British Museum (Natural History), London, 61 p.
- Muspratt, J. 1955. Research on South African Culicini (Diptera, Culicidae). III. A check-list of the species and their distribution, with notes on taxonomy, bionomics and identification. J. ent. Soc. S. Africa 18(2): 149-207.




FIG. 1. Dorsoventral outline of Aedes (Diceromyia) furcifer egg illustrating shape and reticulation. Dorsal aspect is towards the top and anterior end is to the left.




FIG. 2. Enlargement of egg chorion reticulation of Aedes (Diceromyia) furcifer from area indicated by the arrow in figure 1.